

A FATOU-BIEBERBACH DOMAIN IN $\mathbb{C}^2$ WHICH IS NOT RUNGE

ERLEND FORNÆSS WOLD

ABSTRACT. Since a paper by J.P. Rosay and W. Rudin from 1988 there has been an open question whether all Fatou-Bieberbach domains are Runge. We give an example of a Fatou-Bieberbach domain Ω in $\mathbb{C}^2$ which is not Runge. The domain Ω provides (yet) a negative answer to a problem of Bremermann.

1. INTRODUCTION

We give a negative answer to the problem, initially posed by J.P. Rosay and W. Rudin in [6] and later in [4], as to whether all Fatou-Bieberbach domains are Runge:

Theorem 1. *There is a Fatou-Bieberbach domain Ω in $\mathbb{C}^* \times \mathbb{C}$ which is Runge in $\mathbb{C}^* \times \mathbb{C}$ but not in $\mathbb{C}^2$.*

A Fatou-Bieberbach domain is a proper subdomain of $\mathbb{C}^n$ which is biholomorphic to $\mathbb{C}^n$, and a domain $\Omega \subset \mathbb{C}^n$ is said to be Runge (in $\mathbb{C}^n$) if any holomorphic function $f \in \mathcal{O}(\Omega)$ can be approximated uniformly on compacts in Ω by polynomials.

It should be noted that although the domain Ω is not Runge it still has the property that the intersection of Ω with any complex line L is simply connected: Let V be a connected component of $\Omega \cap L$, let $\Gamma \subset V$ be a simple closed curve, and let D denote the disk in L bounded by Γ . Since Γ is null-homotopic in Ω we have that D is contained in $\mathbb{C}^* \times \mathbb{C}$ and so the claim follows from the fact that Ω is Runge in $\mathbb{C}^* \times \mathbb{C}$. Intersecting Ω with a suitable bounded subset of $\mathbb{C}^2$ this gives a negative answer to the problem of Bremermann: "Suppose that D is a Stein domain in $\mathbb{C}^n$ such that for every complex line l in $\mathbb{C}^n$, $l \setminus D$ is connected. Is it true that D is Runge in $\mathbb{C}^n$?" Negative answers to this problem have also recently been given in [1] and [5]. One can in fact show, using an argument as above together with the argument principle, that if $\mathcal{R}$ is a smoothly bounded planar domain and if $\varphi(\mathcal{R})$ is a holomorphic embedding of $\mathcal{R}$ into $\mathbb{C}^2$ with $\varphi(\partial\mathcal{R}) \subset \Omega$, then $\varphi(\mathcal{R}) \subset \Omega$.

The idea of the proof is the following: Observe first that if Ω is a Fatou-Bieberbach domain in $\mathbb{C}^2$ which is Runge, then Ω has the property that if $Y \subset \Omega$ is compact then

Date: January 14, 2007.

2000 Mathematics Subject Classification. 32E20, 32E30, 32H02.

Key words and phrases. Fatou-Bieberbach domains, polynomial convexity, holomorphic maps.

Supported by Schweizerische Nationalfonds grant 200021-116165/1.

its polynomially convex hull

$$\widehat{Y} := \{(z, w) \in \mathbb{C}^2; |P(z, w)| \leq \|P\|_Y \ \forall P \in \mathcal{P}(\mathbb{C}^2)\}$$

is contained in Ω . To prove the theorem we will construct a domain Ω such that $\widehat{Y} \setminus \Omega \neq \emptyset$ for a certain compact set Y . For a compact subset $Y \subset \mathbb{C}^* \times \mathbb{C}$ let $\widehat{Y}_*$ denote the set

$$\widehat{Y}_* := \{(z, w) \in \mathbb{C}^2; |P(z, w)| \leq \|P\|_Y \ \forall P \in \mathcal{O}(\mathbb{C}^* \times \mathbb{C})\}.$$

We say that the set Y is holomorphically convex if $\widehat{Y}_* = Y$. We will first construct (a construction by Stolzenberg) a holomorphically convex compact set $Y \subset \mathbb{C}^* \times \mathbb{C}$ having the property that $\widehat{Y} \cap (\{0\} \times \mathbb{C}) \neq \emptyset$. Y is the disjoint union of two disks in $\mathbb{C}^* \times \mathbb{C}$. We will then use the fact that $\mathbb{C}^* \times \mathbb{C}$ has the *density property* to construct a Fatou-Bieberbach domain $\Omega \subset \mathbb{C}^* \times \mathbb{C}$ such that $Y \subset \Omega$. The domain Ω cannot be Runge.

A few words about the density property and approximation by automorphisms. As defined in [9], *a complex manifold M is said to have the density property if every holomorphic vector field on M can be approximated locally uniformly by Lie combinations of complete vector fields on M* . It was proved in [9] that $\mathbb{C}^* \times \mathbb{C}$ has the density property. In Andersén-Lempert theory the density property corresponds to the fact that in $\mathbb{C}^n$ every entire vector field can be approximated by sums of complete vector fields. This has been studied also in [10].

Using the density property of $\mathbb{C}^* \times \mathbb{C}$ one gets as in [4] (by copying their arguments): Let Ω be an open set in $\mathbb{C}^* \times \mathbb{C}$. For every $t \in [0, 1]$, let φ_t be a biholomorphic map from Ω into $\mathbb{C}^* \times \mathbb{C}$, of class $\mathcal{C}^2$ in $(t, z) \in [0, 1] \times \Omega$. Assume that $\varphi_0 = Id$, and assume that each domain $\Omega_t = \varphi_t(\Omega)$ is Runge in $\mathbb{C}^* \times \mathbb{C}$. Then for every $t \in [0, 1]$ the map φ_t can be approximated on Ω by holomorphic automorphisms of $\mathbb{C}^* \times \mathbb{C}$. In the proof of Theorem 1 we will construct such an isotopy.

We will let π denote the projection onto the first coordinate in $\mathbb{C}^* \times \mathbb{C}$ and in $\mathbb{C}^2$, and we will let $B_\varepsilon(p)$ denote the open ball of radius ε centered at a point p .

2. CONSTRUCTION OF THE SET Y

We start by defining a certain rationally convex subset Y of $\mathbb{C}^2$. The set will be a union of two disjoint polynomially convex disks in $\mathbb{C}^* \times \mathbb{C}$, but the polynomial hull of the union will contain the origin. This construction is taken from [8], page 392-396, and is due to Stolzenberg [7].

Let Ω_1 and Ω_2 be simply connected domains in $\mathbb{C}$, as in Fig.1. below, with smooth boundary, such that if $I_+ = [1, \sqrt{3}]$, $I_- = [-\sqrt{3}, -1]$, then $I_+ \subset \partial\Omega_1$, $I_- \subset \partial\Omega_2$. Require that $\partial\Omega_1$ and $\partial\Omega_2$ meet only twice, that $I_- \subset \Omega_1$, $I_+ \subset \Omega_2$, and, finally, that $\partial\Omega_1 \cup \partial\Omega_2$ be the union of the boundary of the unbounded component of $\mathbb{C} \setminus (\partial\Omega_1 \cup \partial\Omega_2)$, together with the boundary of the component of this set that contains the origin. Let the intersections of the boundaries be the points i and $-i$.




FIG.1. Two smoothly bounded simply connected domains Ω_j with $\partial\Omega_1 \cap \partial\Omega_2 = \pm i$.

We define

$$V_1 = \{(z, w) \in \mathbb{C}^2; z^2 - w \text{ is real and lies in } [0, 1]\},$$

$$V_2 = \{(z, w) \in \mathbb{C}^2; w \text{ is real and lies in } [1, 2]\},$$

$$X_1 = \{(z, w) \in V_1; z \in \partial\Omega_2\},$$

$$X_2 = \{(z, w) \in V_2; z \in \partial\Omega_1\},$$

Note that X_1 and X_2 are totally real annuli, that they are disjoint, and that the origin is contained in the polynomial hull of X_1 . Next we want to remove pieces from X_1 and X_2 to create two disks.

Define

$$\tilde{V}_1 = V_1 \cap \pi^{-1}(I_+),$$

$$\tilde{V}_2 = V_2 \cap \pi^{-1}(I_-),$$

$$Y_1 = \overline{X_1 \setminus \tilde{V}_2},$$

$$Y_2 = \overline{X_2 \setminus \tilde{V}_1}.$$

The set Y will be defined as $Y = Y_1 \cup Y_2$. Note that

$$(*) \quad \tilde{V}_1 \subset \hat{X}_1, \tilde{V}_2 \subset \hat{X}_2.$$

Let us describe what Y_1 and Y_2 looks like over I_- and I_+ respectively. By the equations we see that these sets are contained in $\mathbb{R}^2$. Let (x, y) denote the real parts of (z, w) .

Over I_- we have that Y_1 is the union of the two sets defined by

$$(a) \quad 2 \leq y \leq x^2 \text{ if } -\sqrt{3} \leq x \leq -\sqrt{2},$$

$$(b) \quad x^2 - 1 \leq y \leq 1 \text{ if } -\sqrt{2} \leq x \leq -1.$$

Over I_+ we have that Y_2 is the union of the sets defined by

$$(c) \quad x^2 \leq y \leq 2 \text{ if } 1 \leq x \leq \sqrt{2},$$

$$(d) \quad 1 \leq y \leq x^2 - 1 \text{ if } \sqrt{2} \leq x \leq \sqrt{3}.$$

From these equations we see that Y_1 and Y_2 are disks.

We have that

$$(**) \quad \hat{Y} \text{ contains the origin}$$

because of the following: We already noted that the origin is contained in $\hat{X}_1$, so the claim follows from $(*)$ and the following simpler version of Lemma 29.31, [8], page 392: Let X_1 and X_2 be disjoint compact sets in $\mathbb{C}^N$, and let S_1 and S_2 be relatively open subsets of X_1 and X_2 respectively such that $S_1 \subset \hat{X}_2, S_2 \subset \hat{X}_1$. Then $\widehat{X_1 \cup X_2} = (X_1 \setminus S_1) \cup (X_2 \setminus S_2)$. The reason for this, which was pointed out by the referee, is simply that neither S_1 nor S_2 can contain peak points for the algebra generated by the polynomials on $X_1 \cup X_2$.

3. PROOF OF THEOREM 1

It is proved in [8] that the set Y is rationally convex, and that the sets Y_j are polynomially convex separately. For our construction we need to know that Y is holomorphically convex, so we prove the following:

3.1. Lemma. *We have that Y is holomorphically convex in $\mathbb{C}^* \times \mathbb{C}$.*

Proof. For $j = 1, 2$, let Y_j^+ and Y_j^- denote the sets $Y_j \cap \{\operatorname{Re}(z) \geq 0\}$ and $Y_j \cap \{\operatorname{Re}(z) \leq 0\}$ respectively. Let $Y^+ = Y_1^+ \cup Y_2^+$ and $Y^- = Y_1^- \cup Y_2^-$.

Observe first that Y^+ and Y^- are polynomially convex separately: Assume to get a contradiction that $\widehat{Y^-}$ contains nontrivial points. In that case there exists a graph $G(f)$ of a bounded holomorphic function defined on the topological disk U bounded by

$\pi(Y^-)$, such that $G(f) \subset \widehat{Y^-}$, and such that $(z, f(z)) \in Y^-$ for a.a. (in terms of radial limits if we regard U as a proper disk) $z \in \pi(Y^-)$ (Theorem 20.2. in [2], page 172, holds by the discussion on page 171 even though the fibers over $\pm i$ are not convex). Then for continuity reasons $\overline{G(f)}$ would have to contain nontrivial points of $\widehat{Y^-}$ in the fibers $\{\pm i\} \times \mathbb{C}$ - but as this clearly cannot be the case, we have our contradiction. The case of Y^+ is similar.

Next assume to get a contradiction that there is a point $(z_0, w_0) \in \widehat{Y_*} \setminus Y$ with $\operatorname{Re}(z_0) < 0$. The function $f(z)$ defined to be $(z+i)(z-i)$ on $\pi(Y^-) \cup \{z_0\}$ and zero on $\pi(Y^+)$ can be uniformly approximated on $\pi(Y) \cup \{z_0\}$ by polynomials in z and $\frac{1}{z}$, and so any representing Jensen measure (see [8] Chapter 2) for the functional $g \mapsto g(z_0, w_0)$ would have to be supported on Y^- . But then the point (z_0, w_0) would have to be in the hull of Y^- which is a contradiction. The corresponding conclusion holds for $\operatorname{Re}(z_0) > 0$.

Finally, Rossi's local maximum principle excludes the possibility of there being non-trivial points in the hull contained in $\{\pm i\} \times \mathbb{C}$. $\square$

3.2. Lemma. *Let $p = (z_0, w_0) \in \mathbb{C}^* \times \mathbb{C}$ and let $\varepsilon > 0$. Then there exists an automorphism ψ of $\mathbb{C}^* \times \mathbb{C}$ such that $\psi(Y) \subset B_\varepsilon(p)$.*

Proof. We need to argue that there exists an isotopy as described in the introduction, and we content ourselves by demonstrating that there exist isotopies mapping Y_1 and Y_2 into separate arbitrarily small balls - the rest is trivial. Let $q_j \in Y_j$ be a point for $j = 1, 2$, and let $\delta > 0$. Since Y_j is a smooth disk there clearly exists a smooth map $f^j : [0, 1] \times Y_j \rightarrow Y_j$ such that for each fixed t the map $f_t^j : Y_j \rightarrow Y_j$ is a smooth diffeomorphism, such that f_0^j is the identity, and such that $f_1^j(Y_j) \subset B_\delta(q_j)$. Since Y_j is totally real there exists, by [3] Corollary 3.2, for each $\varepsilon > 0$ a real analytic map $\Phi^j : [0, 1] \times \mathbb{C}^2 \rightarrow \mathbb{C}^2$ such that $\Phi_t^j \in \operatorname{Aut}_{\text{hol}}(\mathbb{C}^2)$ for each t , Φ_0^j is the identity, and $\|f^j - \Phi^j\|_{[0,1] \times Y_j} < \varepsilon$. For small enough ε we restrict Φ^j to a sufficiently small Runge neighborhood of Y_j . $\square$

Proof of Theorem 1: Let G be an automorphism of $\mathbb{C}^* \times \mathbb{C}$ with an attracting fixed point $p \in \mathbb{C}^* \times \mathbb{C}$. It is well known that the basin of attraction of the point p is a Fatou-Bieberbach domain. This domain is clearly contained in $\mathbb{C}^* \times \mathbb{C}$. Denote this domain by $\Omega(G)$. Let ε be a positive real number such that $B_\varepsilon(p) \subset \Omega(G)$. By Lemma 3.2 there is an automorphism ψ of $\mathbb{C}^* \times \mathbb{C}$ such that $\psi(Y) \subset B_\varepsilon(p)$. Then $Y \subset \psi^{-1}(\Omega(G))$. The set $\psi^{-1}(\Omega(G))$ is biholomorphic to $\mathbb{C}^2$, and from (**) in Section 2 we have that $\widehat{Y}$ contains the origin. On the other hand it is clear that $\Omega(G)$ is Runge in $\mathbb{C}^* \times \mathbb{C}$, and so $\psi^{-1}(\Omega(G))$ is Runge in $\mathbb{C}^* \times \mathbb{C}$. $\square$

Acknowledgements. The author would like to thank the referee for comments; in particular for suggesting a simplified proof of Lemma 3.1.

REFERENCES

1. M.Abe: Polynomial Convexity and Strong Disk Property *J. Math. Anal. Appl.* **321**(2006), 32–36.
2. H. Alexander, J. Wermer: Several Complex Variables and Banach Algebras, 3rd ed., *Springer-Verlag New York, Inc.* (1998).
3. F. Forstnerič: Approximation by automorphisms on smooth submanifolds of $\mathbb{C}^n$. *Math. Ann.* **300**(1994), 719–738.
4. F. Forstnerič, J.P. Rosay: Approximation of Biholomorphic Mappings by Automorphisms of $\mathbb{C}^n$. *Invent. Math.* **112** (1993), 112–123.
5. C. Joița: On A Problem of Bremermann Concerning Runge Domains. *Math. Ann.*, **337** , (2007), 395–400.
6. J.P. Rosay. W. Rudin: Holomorphic maps from $\mathbb{C}^n$ to $\mathbb{C}^n$. *Trans. Amer. Mat. Soc.* **310** (1988), 47–86.
7. G. Stolzenberg: On the analytic part of a Runge hull. *Math. Ann.* **164** (1966), 286–290
8. E.L. Stout: The Theorey of Uniform Algebras. *Bogden and Quigly, Inc.* (1971).
9. D. Varolin: The Density Property for Complex Manifolds and Geometric Structures. *J. Geom. Anal.* **11** (2001), 135–160
10. D. Varolin: The Density Property for Complex Manifolds and Geometric Structures II. *Internat. J. Math.* **11** (2000), 837–847.

MATEMATISK INSTITUTT, UNIVERSITETET I OSLO, POSTBOKS 1053 BLINDERN, NO-0316 OSLO, NORWAY

E-mail address: `erlendfw@math.uio.no`